

Training clinicians to proactively ask patients about domestic violence and abuse (DVA) is feasible for sexual health clinics to implement and could increase referrals to specialist services, according to a joint CLAHRC North Thames/CLAHRC West study led by researchers at Queen Mary University of London (QMUL) and the University of Bristol involving over 4,300 women.



The risk of gynaecological and sexual health problems (including sexually transmitted infections, painful sex, vaginal bleeding and recurrent urinary tract infections) is three-fold higher in women who have suffered DVA. Meanwhile, 47 per cent of women attending sexual health services will have experienced DVA at some point in their lives.

Sexual health services can be the first point of contact for women who have experienced DVA, and were listed by the National Institute for Health and Care Excellence (NICE) as a setting in which all patients should be asked about DVA. However, most sexual health professionals have had minimal training in identifying and responding to DVA.

The study looked at the feasibility of sexual health clinics adopting a programme called IRIS

(Identification and Referral to Improve Safety) – a DVA training and referral programme endorsed by NICE, the Royal College of GPs and Department of Health, originally aimed at encouraging GPs to ask patients whether they are experiencing DVA and to make referrals to specialist domestic violence services.



Health Services Research
Short Report

Improving the healthcare response to domestic violence and abuse in sexual health clinics: feasibility study of a training, support and referral intervention 

Alex Hardip Sohal¹, Neha Pathak², Sarah Blake³, Vanessa Apea⁴, Judith Berry⁵, Jayne Bailey^{3, 6}, Chris Griffiths⁷, Gene Feder⁸

Author affiliations 

Abstract

Objectives Sexual health and gynaecological problems are the most consistent and largest physical health differences between abused and non-abused female populations. Sexual health services are well placed to identify and support patients experiencing domestic violence and abuse (DVA). Most sexual health professionals have had minimal DVA training despite English National Institute for Health and Care Excellence recommendations. We sought to determine the feasibility of an evidence-based complex DVA training intervention in female sexual health walk-in services (IRIS ADVISE: Identification and Referral to Improve Safety whilst Assessing Domestic Violence in Sexual Health Environments).

Methods An adaptive mixed method pilot study in the female walk-in service of two sexual health clinics. Following implementation and evaluation at site 1, the intervention was refined before implementation at site 2. The intervention comprised electronic prompts, multidisciplinary training sessions, clinic materials and simple referral pathways to IRIS ADVISE advocate-educators (AEs). The pilot lasted 7 weeks at site 1 and 12 weeks at site 2. Feasibility outcomes were to assign a supportive DVA clinical lead, an IRIS ADVISE AE employed by a local DVA service provider, adapt electronic records, develop local referral pathways, assess whether enquiry, identification and referral rates were measurable.

Results Both sites achieved all feasibility outcomes: appointing a supportive DVA clinical lead and IRIS ADVISE AE, establishing links with a local DVA provider, adapting electronic records, developing local referral pathways and rates of enquiry, identification and referral were found to be measurable. Site 1: 10% enquiry rate (n=267), 4% identification rate (n=16) and eight AE referrals. Site 2: 61% enquiry rate (n=1090), a 7% identification rate (n=79) and eight AE referrals.

Conclusions IRIS ADVISE can be successfully developed and implemented in sexual health clinics. It fulfils the unmet need for DVA training. Longer-term evaluation is recommended.

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited.

The team tested the intervention in two female walk-in sexual health services; an east London clinic serving an inner-city multi-ethnic population, and a Bristol clinic serving an urban population.

In the east London clinic over seven weeks, 267 out of 2,568 women attending were asked about DVA, with 16 of those (6 per cent) saying that they were affected by abuse. Overall, eight of the women affected by abuse (50 per cent) were referred to specialist services.

In the Bristol clinic over twelve weeks, 1,090 out of 1,775 women attending were asked about

DVA, with 79 of those (7 per cent) saying they were affected by abuse. Overall, eight of the women affected by abuse (10 per cent), were referred to the specialist services.

Lead CLAHRC North Thames researcher Dr Alex Sohal (pictured left) said:

“Women attend sexual health clinics for care of their sexual health but little thought is given to whether the relationship with the person that a woman has sex with directly harms her health. Without training, system level changes and senior managerial support, clinicians end up ignoring DVA in consultations or have an arbitrary approach that fails many women affected by DVA. Not only is this a feasible intervention for a sexual health clinic setting, but we also found that clinical leads and busy local DVA service providers were incredibly supportive, with many people understanding the importance of making this work.”



.”

Read the full paper:

[Improving the healthcare response to domestic violence and abuse in sexual health clinics: feasibility study of a training, support and referral intervention](#)

Sohal AH, Pathak N, Blake S, et al. Sex Transm Infect Published Online First: doi:10.1136/sextrans-2016-052866